

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22201

Important Instructions to Examiners:

- 1) The answers should be examined by key words and not as word-to-word as given in the model answer scheme.
- 2) The model answer and the answer written by candidate may vary but the examiner may try to assess the understanding level of the candidate.
- 3) The language errors such as grammatical, spelling errors should not be given more importance (Not applicable for subject English and Communication Skills).
- 4) While assessing figures, examiner may give credit for principal components indicated in the figure. The figures drawn by candidate and model answer may vary. The examiner may give credit for any equivalent figure drawn.
- 5) Credits may be given step wise for numerical problems. In some cases, the assumed constant values may vary and there may be some difference in the candidate's answer and model answer.
- 6) In case of some questions credit may be given by judgement on part of examiner of relevant answer based on candidate's understanding.
- 7) For programming language papers, credit may be given to any other program based on equivalent concept.

Q. No.	Sub Q. N.	Answer	Marking Scheme	
1.		Attempt any FIVE of following:	10	
	a)	If $f(x) = 16^x - \log_2 x$ find $f\left(\frac{1}{4}\right)$	02	
	Ans	$f(x) = 16^x - \log_2 x$		
		$f\left(\frac{1}{4}\right) = 16^{\frac{1}{4}} - \log_2\left(\frac{1}{4}\right)$	1	
		$= 2$	1	

	b)	If $f(x) = ax^2 - bx - 1, f(2) = 5, f(-2) = 10$ find a and b.	02	
	Ans	$f(x) = ax^2 - bx - 1$		
		$f(2) = 5$		
		$4a - 2b - 1 = 5$		
	$4a - 2b = 6$ -----(1)	$\frac{1}{2}$		
	$f(-2) = 10$			
	$4a + 2b - 1 = 10$			
	$4a + 2b = 11$ -----(2)	$\frac{1}{2}$		
	From (1) and (2)	$\frac{1}{2}$		
	$a = \frac{17}{8}$	$\frac{1}{2}$		
	$b = \frac{5}{4}$	$\frac{1}{2}$		

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
1.	b)	Find $\frac{dy}{dx}$, if $y = x \sin^{-1} x$	02
	Ans	$y = x \sin^{-1} x$ $\frac{dy}{dx} = x \frac{1}{\sqrt{1-x^2}} + \sin^{-1} x \cdot 1$ $= \frac{x}{\sqrt{1-x^2}} + \sin^{-1} x$	2
	c)	Evaluate: $\int \frac{dx}{3x^2+4}$	02
Ans	$\int \frac{dx}{3x^2+4}$ $= \int \frac{dx}{(\sqrt{3}x)^2 + 2^2}$ $= \frac{1}{2} \tan^{-1} \left(\frac{\sqrt{3}x}{2} \right) \frac{1}{\sqrt{3}} + c$ $= \frac{1}{2\sqrt{3}} \tan^{-1} \left(\frac{\sqrt{3}x}{2} \right) + c$	<p>1/2</p> <p>1</p> <p>1/2</p>	
e)	Evaluate $\int \sin^3 x dx$	02	
Ans	$\int \sin^3 x dx$ <p>since $\sin 3x = 3 \sin x - 4 \sin^3 x \quad \therefore \sin^3 x = \frac{1}{4}(3 \sin x - \sin 3x)$</p> $\therefore \int \frac{1}{4}(3 \sin x - \sin 3x) dx$ $= \frac{1}{4} \left(-3 \cos x + \frac{\cos 3x}{3} \right) + c$ <p>OR</p> $\int \sin^3 x dx$ $= \int \sin^2 x \sin x dx$ $= \int (1 - \cos^2 x) \sin x dx$ <p>Put $\cos x = t$</p> $\therefore -\sin x dx = dt$ $\therefore \sin x dx = -dt$	<p>1</p> <p>1</p> <p>1/2</p> <p>1/2</p>	

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
1.	e)	$\therefore \int (1-t^2)(-dt)$ $= -\int (1-t^2)dt$ $= -\left(t - \frac{t^3}{3}\right) + c$ $= -\left(\cos x - \frac{\cos^3 x}{3}\right) + c$	<p>1/2</p> <p>1/2</p>
	f)	<p>Find the volume obtained by revolving the area under the curve $9x^2 - 4y^2 = 36$ in the interval from $x = 2$ to $x = 4$ about x-axis</p> <p>Ans $9x^2 - 4y^2 = 36$</p> $y^2 = \frac{9}{4}(x^2 - 4)$ $\text{volume} = \pi \int_a^b y^2 dx$ $= \pi \int_2^4 \frac{9}{4}(x^2 - 4) dx$ $= \frac{9\pi}{4} \left[\frac{x^3}{3} - 4x \right]_2^4$ $= \frac{9\pi}{4} \left[\left(\frac{4^3}{3} - 4(4) \right) - \left(\frac{2^3}{3} - 4(2) \right) \right]$ $= 24\pi$	<p>02</p> <p>1/2</p> <p>1</p> <p>1/2</p>
	g)	<p>Find order and degree of the differential equation $\frac{d^2y}{dx^2} = \left(y + \frac{dy}{dx}\right)^{\frac{3}{2}}$</p> <p>Ans $\frac{d^2y}{dx^2} = \left(y + \frac{dy}{dx}\right)^{\frac{3}{2}}$</p> <p>Squaring on both sides</p> $\left(\frac{d^2y}{dx^2}\right)^2 = \left(y + \frac{dy}{dx}\right)^3$ <p>\therefore Order = 2</p> <p>\therefore Degree = 2</p>	<p>02</p> <p>1</p> <p>1</p>
2.		<p>Attempt any THREE of the following:</p>	12

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
2.	a)	<p>If $x^p y^q = (x + y)^{p+q}$ show that $\frac{dy}{dx} = \frac{y}{x}$</p> <p>Ans $x^p y^q = (x + y)^{p+q}$</p> $\log(x^p y^q) = \log(x + y)^{p+q}$ $\log x^p + \log y^q = (p + q) \log(x + y)$ $p \log x + q \log y = (p + q) \log(x + y)$ $p \frac{1}{x} + q \frac{1}{y} \frac{dy}{dx} = (p + q) \left(\frac{1}{x + y} \left(1 + \frac{dy}{dx} \right) \right)$ $\frac{p}{x} + \frac{q}{y} \frac{dy}{dx} = \frac{p + q}{x + y} + \frac{p + q}{x + y} \frac{dy}{dx}$ $\frac{q}{y} \frac{dy}{dx} - \frac{p + q}{x + y} \frac{dy}{dx} = \frac{p + q}{x + y} - \frac{p}{x}$ $\frac{dy}{dx} \left(\frac{q}{y} - \frac{p + q}{x + y} \right) = \frac{p + q}{x + y} - \frac{p}{x}$ $\frac{dy}{dx} \left(\frac{qx + qy - py - qy}{y(x + y)} \right) = \frac{px + qx - px - py}{x(x + y)}$ $\frac{dy}{dx} \left(\frac{qx - py}{y} \right) = \frac{qx - py}{x}$ $\frac{dy}{dx} = \frac{y}{x}$	<p>04</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p>
	b)	<p>If $y = 3 \sin \theta - 2 \sin^3 \theta$, $x = 3 \cos \theta - 2 \cos^3 \theta$ find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{4}$</p> <p>Ans $y = 3 \sin \theta - 2 \sin^3 \theta$</p> $\therefore \frac{dy}{d\theta} = 3 \cos \theta - 6 \sin^2 \theta \cdot \cos \theta$ $= 3 \cos \theta (1 - 2 \sin^2 \theta)$ $x = 3 \cos \theta - 2 \cos^3 \theta$ $\frac{dx}{d\theta} = -3 \sin \theta + 6 \cos^2 \theta \cdot \sin \theta$ $= -3 \sin \theta (1 - 2 \cos^2 \theta)$ $\therefore \frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{3 \cos \theta (1 - 2 \sin^2 \theta)}{-3 \sin \theta (1 - 2 \cos^2 \theta)}$ $= \frac{3 \cos \theta (\cos 2\theta)}{-3 \sin \theta (-\cos 2\theta)}$	<p>04</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p>

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228

Page No.04/19

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code **22201**

Q. No.	Sub Q. N.	Answer	Marking Scheme
2.	b)	$= \cot \theta$ $\therefore \text{at } \theta = \frac{\pi}{4}$ $\therefore \frac{dy}{dx} = \cot \frac{\pi}{4}$ $= 1$	½
	c)	<p>Find the radius of curvature of the curve $xy = c$ at point (c, c)</p> <p>Ans $xy = c$</p> $x \frac{dy}{dx} + y \cdot 1 = 0$ $\frac{dy}{dx} = -\frac{y}{x}$ $\frac{d^2y}{dx^2} = -\left[\frac{x \frac{dy}{dx} - y}{x^2} \right]$ <p>at point (c, c)</p> $\frac{dy}{dx} = -\frac{c}{c} = -1$ $\frac{d^2y}{dx^2} = -\frac{[c(-1) - c]}{c^2}$ $= \frac{2}{c}$ $\therefore \text{radius of curvature} = \frac{\left(1 + \left(\frac{dy}{dx} \right)^2 \right)^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$ $= \frac{(1 + (-1)^2)^{\frac{3}{2}}}{\frac{2}{c}}$ $= 2^{\frac{3}{2}} c \text{ or } \sqrt{2}c$	04
	d)	<p>Discuss the maxima and minima of the function "$\tan x - 2x$"</p> <p>Ans Let $y = \tan x - 2x$</p>	04

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
2.	d)	$\therefore \frac{dy}{dx} = \sec^2 x - 2$ $\therefore \frac{d^2y}{dx^2} = 2 \sec x \sec x \tan x = 2 \sec^2 x \tan x$ <p>consider $\frac{dy}{dx} = 0$</p> $\therefore \sec^2 x - 2 = 0$ $\therefore \sec^2 x = 2$ $\therefore \sec x = \sqrt{2}, -\sqrt{2}$ $\therefore x = \frac{\pi}{4}, \frac{3\pi}{4}$ <p>at $x = \frac{\pi}{4}$</p> $\therefore \frac{d^2y}{dx^2} = 2 \sec^2 \frac{\pi}{4} \tan \frac{\pi}{4} = 2(2)(1) = 4 > 0$ <p>\therefore function is minimum at $x = \frac{\pi}{4}$</p> $\therefore y_{\min} = \tan \frac{\pi}{4} - 2 \left(\frac{\pi}{4} \right) = 1 - \frac{\pi}{2}$ <p>at $x = \frac{3\pi}{4}$</p> $\therefore \frac{d^2y}{dx^2} = 2 \sec^2 \left(\frac{3\pi}{4} \right) \tan \left(\frac{3\pi}{4} \right) = 2(2)(-1) = -4 < 0$ <p>\therefore function is maximum at $x = \frac{3\pi}{4}$</p> $\therefore y_{\max} = \tan \left(\frac{3\pi}{4} \right) - 2 \left(\frac{3\pi}{4} \right) = -1 - \frac{3\pi}{2}$	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>
3.		<p>Attempt any THREE of the following:</p> <p>a) Find the equation of tangent and normal to the curve $y = x(2-x)$ at point $(2,0)$</p> <p>Ans $y = x(2-x)$</p> $\therefore y = 2x - x^2$ $\therefore \frac{dy}{dx} = 2 - 2x$ <p>at $(2,0)$</p> $\therefore \frac{dy}{dx} = 2 - 2(2)$	<p>12</p> <p>04</p> <p>1</p>

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
3.	a)	$\therefore \frac{dy}{dx} = -2$ $\therefore \text{slope of tangent, } m = -2$ <p>Equation of tangent at (2,0) is</p> $y - 0 = -2(x - 2)$ $\therefore y = -2x + 4$ $\therefore 2x + y - 4 = 0$ $\therefore \text{slope of normal, } m' = \frac{-1}{m} = \frac{1}{2}$ <p>Equation of normal at (2,0) is</p> $y - 0 = \frac{1}{2}(x - 2)$ $\therefore 2y = x - 2$ $\therefore x - 2y - 2 = 0$	<p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p>
	b)	<p>Find $\frac{dy}{dx}$, $y = (\sin^{-1} x)^x + (\cos x)^{\sin x}$</p> <p>Ans Let $u = (\sin^{-1} x)^x$</p> $\therefore \log u = \log (\sin^{-1} x)^x$ $\therefore \log u = x \log (\sin^{-1} x)$ $\therefore \frac{1}{u} \frac{du}{dx} = x \cdot \frac{1}{\sin^{-1} x} \cdot \frac{1}{\sqrt{1-x^2}} + \log (\sin^{-1} x) \cdot 1$ $\therefore \frac{1}{u} \frac{du}{dx} = \frac{x}{\sqrt{1-x^2} \sin^{-1} x} + \log (\sin^{-1} x)$ $\therefore \frac{du}{dx} = u \left(\frac{x}{\sqrt{1-x^2} \sin^{-1} x} + \log (\sin^{-1} x) \right)$ $\therefore \frac{du}{dx} = (\sin^{-1} x)^x \left(\frac{x}{\sqrt{1-x^2} \sin^{-1} x} + \log (\sin^{-1} x) \right)$ <p>Let $v = (\cos x)^{\sin x}$</p> $\therefore \log v = \log (\cos x)^{\sin x}$ $\therefore \log v = \sin x \log (\cos x)$ $\therefore \frac{1}{v} \frac{dv}{dx} = \sin x \cdot \frac{1}{\cos x} \cdot (-\sin x) + \log (\cos x) \cdot \cos x$	<p>04</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p>

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
3.	b)	$\therefore \frac{1}{v} \frac{dv}{dx} = -\sin x \tan x + \log(\cos x) \cdot \cos x$ $\therefore \frac{dv}{dx} = v[-\sin x \tan x + \log(\cos x) \cdot \cos x]$ $\therefore \frac{dv}{dx} = (\cos x)^{\sin x} [-\sin x \tan x + \log(\cos x) \cdot \cos x]$ $\therefore \frac{dy}{dx} = (\sin^{-1} x)^x \left(\frac{x}{\sqrt{1-x^2} \sin^{-1} x} + \log(\sin^{-1} x) \right) + (\cos x)^{\sin x} [-\sin x \tan x + \log(\cos x) \cdot \cos x]$	<p>1/2</p> <p>1/2</p>
	c)	<p>If $y = \tan^{-1} \left[\frac{5x-4}{5+4x} \right]$ find $\frac{dy}{dx}$</p> <p>Ans $y = \tan^{-1} \left[\frac{5x-4}{5+4x} \right]$</p> $y = \tan^{-1} \left[\frac{x - \frac{4}{5}}{1 + \frac{4}{5}x} \right]$ $y = \tan^{-1} x - \tan^{-1} \frac{4}{5}$ $\frac{dy}{dx} = \frac{1}{1+x^2} - 0$ $\frac{dy}{dx} = \frac{1}{1+x^2}$	<p>04</p> <p>1</p> <p>2</p> <p>1</p>
	d)	<p>Evaluate $\int \frac{\sec^2 x}{(1+\tan x)(2+\tan x)} dx$</p> <p>Ans $\int \frac{\sec^2 x}{(1+\tan x)(2+\tan x)} dx$</p> <p>Let $\tan x = t$</p> $\therefore \sec^2 x dx = dt$ $= \int \frac{1}{(1+t)(2+t)} dt$ <p>Consider</p> $\frac{1}{(1+t)(2+t)} = \frac{A}{1+t} + \frac{B}{2+t}$	<p>04</p> <p>1/2</p>

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
	d)	$\therefore 1 = A(2+t) + B(1+t)$ $\text{Put } t = -1 \quad \therefore A = 1$ $\text{Put } t = -2 \quad \therefore B = -1$ $\therefore \frac{1}{(1+t)(2+t)} = \frac{1}{1+t} + \frac{-1}{2+t}$ $\therefore \int \frac{1}{(1+t)(2+t)} dt = \int \left(\frac{1}{1+t} + \frac{-1}{2+t} \right) dt$ $= 1 \log(1+t) - 1 \log(2+t) + c$ $= \log \left(\frac{1+t}{2+t} \right) + c$ $= \log \left(\frac{1 + \tan x}{2 + \tan x} \right) + c$ <p>OR</p> $\int \frac{\sec^2 x}{(1 + \tan x)(2 + \tan x)} dx$ $\text{Put } \tan x = t$ $\therefore \sec^2 x dx = dt$ $\int \frac{1}{(1+t)(2+t)} dt$ $= \int \frac{1}{t^2 + 3t + 2} dt$ $\text{Third Term} = \frac{3^2}{4} = \frac{9}{4}$ $= \int \frac{1}{t^2 + 4t + \frac{9}{4} - \frac{9}{4} + 2} dt$ $= \int \frac{1}{\left(t + \frac{3}{2}\right)^2 - \left(\frac{1}{2}\right)^2} dt$ $= \frac{1}{2 \cdot \frac{1}{2}} \log \left \frac{t + \frac{3}{2} - \frac{1}{2}}{t + \frac{3}{2} + \frac{1}{2}} \right + c$ $= \log \left \frac{t+1}{t+2} \right + c$ $= \log \left \frac{\tan x + 1}{\tan x + 2} \right + c$	<p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p>
4.		<p>Attempt any THREE of the following:</p>	12
	a)	<p>Evaluate : $\int \frac{1}{2x^2 + 3x + 1} dx$</p>	04

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
4.	a)Ans	$\int \frac{1}{2x^2 + 3x + 1} dx = \frac{1}{2} \int \frac{1}{x^2 + \frac{3}{2}x + \frac{1}{2}} dx$ $\text{Third term} = \left(\frac{1}{2} \times \frac{3}{2}\right)^2 = \frac{9}{16}$ $= \frac{1}{2} \int \frac{1}{x^2 + \frac{3}{2}x + \frac{9}{16} - \frac{9}{16} + \frac{1}{2}} dx$ $= \frac{1}{2} \int \frac{1}{\left(x + \frac{3}{4}\right)^2 - \left(\frac{1}{4}\right)^2} dx$ $= \frac{1}{2} \left[\frac{1}{2\left(\frac{1}{4}\right)} \log \left(\frac{x + \frac{3}{4} - \frac{1}{4}}{x + \frac{3}{4} + \frac{1}{4}} \right) \right] + c$ $= \log \left(\frac{2x+1}{2x+2} \right) + c$ <p style="text-align: center;">OR</p> $\int \frac{1}{2x^2 + 3x + 1} dx = \int \frac{1}{(2x+1)(x+1)} dx$ <p>Let $\frac{1}{(2x+1)(x+1)} = \frac{A}{2x+1} + \frac{B}{x+1}$</p> $1 = A(x+1) + B(2x+1)$ <p>Put $x = \frac{-1}{2}$</p> $\therefore A = 2$ <p>Put $x = -1$</p> $\therefore B = -1$ $\frac{1}{(2x+1)(x+1)} = \frac{2}{2x+1} + \frac{-1}{x+1}$ $\int \frac{1}{(2x+1)(x+1)} dx = \int \left(\frac{2}{2x+1} + \frac{-1}{x+1} \right) dx$ $= \frac{2 \log(2x+1)}{2} - \log(x+1) + c$ $= \log(2x+1) - \log(x+1) + c$ <p>OR</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1</p>

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
4.	a)	$\int \frac{1}{2x^2 + 3x + 1} dx$	
	Ans	<p>Third term = $\frac{(M.T.)^2}{4(F.T.)} = \frac{9}{8}$</p> $= \int \frac{1}{2x^2 + 3x + \frac{9}{8} - \frac{9}{8} + 1} dx$ $= \int \frac{1}{\left(\sqrt{2x} + \frac{3}{\sqrt{8}}\right)^2 - \frac{1}{8}} dx$ $= \int \frac{1}{\left(\sqrt{2x} + \frac{3}{\sqrt{8}}\right)^2 - \left(\frac{1}{\sqrt{8}}\right)^2} dx$ $= \frac{1}{\sqrt{2}} \left[\frac{1}{2\left(\frac{1}{\sqrt{8}}\right)} \log \left(\frac{\sqrt{2x} + \frac{3}{\sqrt{8}} - \frac{1}{\sqrt{8}}}{\sqrt{2x} + \frac{3}{\sqrt{8}} + \frac{1}{\sqrt{8}}} \right) \right] + c$ $= \log \left(\frac{2x+1}{2x+2} \right) + c$	1 1 1 1 1
	b)	<p>Evaluate $\int \frac{dx}{1 + \sin x + \cos x}$</p>	04
	Ans	$\int \frac{dx}{1 + \sin x + \cos x}$ <p>Put $\tan \frac{x}{2} = t \quad \therefore \sin x = \frac{2t}{1+t^2}, \cos x = \frac{1-t^2}{1+t^2}, dx = \frac{2dt}{1+t^2}$</p> $\therefore \int \frac{dx}{1 + \sin x + \cos x} = \int \frac{1}{1 + \frac{2t}{1+t^2} + \left(\frac{1-t^2}{1+t^2}\right)} \cdot \frac{2dt}{1+t^2}$ $= \int \frac{2 dt}{1+t^2 + 2t + 1 - t^2} dt$ $= 2 \int \frac{1}{2t+2} dt$ $= \int \frac{dt}{t+1}$ $= \log(t+1) + c$ $= \log \left(\tan \frac{x}{2} + 1 \right) + c$	1 1 1 1

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
4.	c)	<p>Evaluate : $\int \frac{x \sin^{-1} x}{\sqrt{1-x^2}} dx$</p> <p>Ans $\int \frac{x \sin^{-1} x}{\sqrt{1-x^2}} dx$ put $\sin^{-1} x = t \quad \therefore x = \sin t$ $\frac{1}{\sqrt{1-x^2}} dx = dt$ $\int t \sin t dt$ $= t \int \sin t dt - \int \left(\int \sin t dt \right) \frac{d}{dt}(t) dt$ $= t(-\cos t) - \int -\cos t \cdot 1 \cdot dt$ $= -t \cos t + \sin t + c$ $= -\sin^{-1} x \cos(\sin^{-1} x) + \sin(\sin^{-1} x) + c$</p>	<p>04</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p>
	d)	<p>Evaluate: $\int_0^{\pi/2} \frac{\tan x}{1 + \tan x} dx$</p> <p>Ans Let $I = \int_0^{\pi/2} \frac{\tan x}{1 + \tan x} dx$ $= \int_0^{\pi/2} \frac{\sin x}{1 + \frac{\sin x}{\cos x}} dx$ $\therefore I = \int_0^{\pi/2} \frac{\sin x}{\sin x + \cos x} dx \quad \text{-----(1)}$ $I = \int_0^{\pi/2} \frac{\sin(\pi/2 - x)}{\sin(\pi/2 - x) + \cos(\pi/2 - x)} dx$ $\therefore I = \int_0^{\pi/2} \frac{\cos x}{\cos x + \sin x} dx \quad \text{-----(2)}$ add (1) and (2) $I + I = \int_0^{\pi/2} \frac{\sin x}{\sin x + \cos x} dx + \int_0^{\pi/2} \frac{\cos x}{\cos x + \sin x} dx$</p>	<p>04</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p>

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
4.	d)	$2I = \int_0^{\pi/2} \frac{\sin x + \cos x}{\sin x + \cos x} dx$ $2I = \int_0^{\pi/2} 1 dx$ $2I = [x]_0^{\pi/2}$ $2I = \frac{\pi}{2} - 0$ $I = \frac{\pi}{4}$	<p>1/2</p> <p>1/2</p> <p>1/2</p>
	e)	<p>Evaluate: $\int \frac{x}{(x^2+4)(x^2+9)} dx$</p> <p>Ans $\int \frac{x}{(x^2+4)(x^2+9)} dx$</p> <p>Put $x^2 = t$</p> <p>$2x dx = dt$</p> <p>$x dx = \frac{dt}{2}$</p> <p>$\int \frac{\frac{dt}{2}}{(t+4)(t+9)}$</p> <p>$= \frac{1}{2} \int \frac{dt}{(t+4)(t+9)}$</p> <p>$\frac{1}{(t+4)(t+9)} = \frac{A}{t+4} + \frac{B}{t+9}$</p> <p>$1 = A(t+9) + B(t+4)$</p> <p>Put $t = -4 \quad \therefore A = \frac{1}{5}$</p> <p>Put $t = -9 \quad \therefore B = -\frac{1}{5}$</p> <p>$\frac{1}{(t+4)(t+9)} = \frac{1}{5} \frac{1}{t+4} - \frac{1}{5} \frac{1}{t+9}$</p> <p>$\int \frac{dt}{(t+4)(t+9)} = \int \left(\frac{1}{5} \frac{1}{t+4} - \frac{1}{5} \frac{1}{t+9} \right) dt$</p>	<p>04</p> <p>1/2</p> <p>1</p> <p>1</p> <p>1/2</p>

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
5.	e)	$= \frac{1}{5} \log(t+4) - \frac{1}{5} \log(t+9) + c$ $= \frac{1}{5} \log(x^2+4) - \frac{1}{5} \log(x^2+9) + c$	<p>½</p> <p>½</p>
		<p>Attempt any TWO of the following:</p> <p>Find area of between the curve $y^2 - 2x = 0$ and $y^2 + 4x - 12 = 0$</p>	12
	a)	$y^2 = 2x$ -----(1)	06
	Ans	$y^2 = 12 - 4x$ $\therefore 2x = 12 - 4x$ $\therefore 6x = 12$ $\therefore x = 2, y = \pm 2$ $\therefore x = \frac{y^2}{2}, x = \frac{12 - y^2}{4}$ Area $A = \int_a^b (x_1 - x_2) dy$ $\therefore A = \int_{-2}^2 \left(\frac{12 - y^2}{4} - \frac{y^2}{2} \right) dy$ $\therefore A = \frac{3}{4} \int_{-2}^2 (4 - y^2) dy$ $\therefore A = \frac{3}{4} \left(4y - \frac{y^3}{3} \right)_{-2}^2$ $\therefore A = \frac{3}{4} \left(4(2) - \frac{(2)^3}{3} - \left(4(-2) - \frac{(-2)^3}{3} \right) \right)$ $\therefore A = 8$	1
	b)	Attempt the following:	06
	(i)	Form the differential equation If $y = A \cos(\log x) + B \sin(\log x)$	03
	Ans	$y = A \cos(\log x) + B \sin(\log x)$ $\therefore \frac{dy}{dx} = -\frac{A \sin(\log x)}{x} + \frac{B \cos(\log x)}{x}$ $\therefore x \frac{dy}{dx} = -A \sin(\log x) + B \cos(\log x)$ $\therefore x \frac{d^2 y}{dx^2} + \frac{dy}{dx} = -\frac{A \cos(\log x)}{x} - \frac{B \sin(\log x)}{x}$ $x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} = -A \cos(\log x) - B \sin(\log x)$	1
			½
			1

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
5.	b)i)	$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} = -y$ $x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + y = 0$	½
	b)ii)	Solve $x \log x \frac{dy}{dx} + y = 2 \log x$	03
	Ans	$x \log x \frac{dy}{dx} + y = 2 \log x$ $\frac{dy}{dx} + \frac{y}{x \log x} = \frac{2 \log x}{x \log x}$ $\frac{dy}{dx} + \frac{y}{x \log x} = \frac{2}{x}$ $I.F = e^{\int \frac{1}{x \log x} dx}$ $= e^{\log(\log x)} = \log x$ <p>Solution is,</p> $y \cdot \log x = \int \frac{2}{x} \cdot \log x dx$ <p>Let $I_1 = \int \frac{2}{x} \cdot \log x dx$</p> <p>Put $\log x = t$</p> $\frac{1}{x} dx = dt$ $\therefore I_1 = 2 \int t dt$ $= 2 \frac{t^2}{2} + c$ $= (\log x)^2 + c$ $y \cdot \log x = (\log x)^2 + c$	½ ½ ½ ½ ½ ½
c)	A circular column of radius 'x' and having depth y support a load. The equation of equilibrium is $2 \frac{dy}{dx} - kx = 0$ where 'k' is constant. Find the relation between x and y.	06	
	Ans	$2 \frac{dy}{dx} - kx = 0$ $2 \frac{dy}{dx} = kx$	

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22201

Q. No.	Sub Q. N.	Answer	Marking Scheme												
5.	c)	$\frac{dy}{dx} = \frac{k}{2}x$ $dy = \frac{k}{2}x dx$ $\int dy = \int \frac{k}{2}x dx$ $y = \frac{k}{2} \frac{x^2}{2} + c_1$ $4y = kx^2 + 4c$ $4y = kx^2 + c$	<p>½</p> <p>1</p> <p>1</p> <p>2</p> <p>1</p> <p>½</p>												
6.		<p>Attempt any TWO of the following:</p> <p>a) Using Simpson's 1/3rd rule evaluate $\int_0^2 \frac{1}{1+x^3} dx$ with $n=4$.</p> <p>Ans Let $y = \frac{1}{1+x^3}$ $a=0, b=2$ and $n=4$</p> $\therefore h = \frac{b-a}{n} = \frac{2-0}{4} = \frac{1}{2}$ <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>0</td> <td>$\frac{1}{2}$</td> <td>1</td> <td>$\frac{3}{2}$</td> <td>2</td> </tr> <tr> <td>$y = \frac{1}{1+x^3}$</td> <td>1</td> <td>$\frac{8}{9}$</td> <td>$\frac{1}{2}$</td> <td>$\frac{8}{35}$</td> <td>$\frac{1}{9}$</td> </tr> </table> <p>Using Simpson's 1/3rd rule</p> $\int_a^b f(x) dx = \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$ $\therefore \int_0^1 f(x) dx = \frac{1/2}{3} \left[\left(1 + \frac{1}{9}\right) + 4\left(\frac{8}{9} + \frac{8}{35}\right) + 2\left(\frac{1}{2}\right) \right]$ $\therefore \int_0^1 \frac{1}{1+x^3} dx = 1.0968$ <p>OR</p> <p>Let $y = \frac{1}{1+x^3}$ $a=0, b=2$ and $n=4$</p> $\therefore h = \frac{b-a}{n} = \frac{2-0}{4} = \frac{1}{2} = 0.5$	x	0	$\frac{1}{2}$	1	$\frac{3}{2}$	2	$y = \frac{1}{1+x^3}$	1	$\frac{8}{9}$	$\frac{1}{2}$	$\frac{8}{35}$	$\frac{1}{9}$	<p>12</p> <p>06</p> <p>1</p> <p>2</p> <p>2</p> <p>1</p> <p>1</p>
x	0	$\frac{1}{2}$	1	$\frac{3}{2}$	2										
$y = \frac{1}{1+x^3}$	1	$\frac{8}{9}$	$\frac{1}{2}$	$\frac{8}{35}$	$\frac{1}{9}$										

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code:

22201

Q. No.	Sub Q. N.	Answer	Marking Scheme																			
6.	a)	<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>0</td> <td>0.5</td> <td>1</td> <td>1.5</td> <td>2</td> </tr> <tr> <td>$y = \frac{1}{1+x^3}$</td> <td>1</td> <td>0.8889</td> <td>0.5</td> <td>0.2286</td> <td>0.1111</td> </tr> </table> <p>Using Simpson's $1/3^{rd}$ rule</p> $\int_a^b f(x) dx = \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$ $\therefore \int_0^1 f(x) dx = \frac{0.5}{3} [(1 + 0.1111) + 4(0.8889 + 0.2286) + 2(0.5)]$ $\int_0^1 \frac{1}{1+x^2} dx = 1.0969$	x	0	0.5	1	1.5	2	$y = \frac{1}{1+x^3}$	1	0.8889	0.5	0.2286	0.1111	2							
	x	0	0.5	1	1.5	2																
	$y = \frac{1}{1+x^3}$	1	0.8889	0.5	0.2286	0.1111																
b)	<p>Using Simpson's $3/8^{th}$ rule, evaluate $\int_0^{\frac{\pi}{2}} \cos x dx$ with $n = 8$</p> <p>Here $n = 8$</p> <p>Ans $y = \cos x \quad a = 0, \quad b = \frac{\pi}{2}$</p> $\therefore h = \frac{b-a}{n} = \frac{\frac{\pi}{2} - 0}{8} = \frac{\pi}{16}$ <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>0</td> <td>$\frac{\pi}{16}$</td> <td>$\frac{\pi}{8}$</td> <td>$\frac{3\pi}{16}$</td> <td>$\frac{\pi}{4}$</td> <td>$\frac{5\pi}{16}$</td> <td>$\frac{3\pi}{8}$</td> <td>$\frac{7\pi}{16}$</td> <td>$\frac{\pi}{2}$</td> </tr> <tr> <td>$y = \cos x$</td> <td>1</td> <td>0.9808</td> <td>0.9239</td> <td>0.8315</td> <td>0.7071</td> <td>0.5556</td> <td>0.3827</td> <td>0.1951</td> <td>0</td> </tr> </table> <p>Using Simpson's $3/8^{th}$ rule.</p> $\int_a^b f(x) dx = \frac{3h}{8} [(y_0 + y_n) + 3(y_1 + y_2 + y_4 + \dots + y_{n-1}) + 2(y_3 + y_6 + \dots + y_{n-3})]$ $\therefore \int_0^{\frac{\pi}{2}} \cos x dx = \frac{3\left(\frac{\pi}{16}\right)}{8} [(1 + 0) + 3(0.9808 + 0.9239 + 0.7071 + 0.5556 + 0.1951) + 2(0.8315 + 0.3827)]$ $\therefore \int_0^{\frac{\pi}{2}} \cos x dx = 0.9952$	x	0	$\frac{\pi}{16}$	$\frac{\pi}{8}$	$\frac{3\pi}{16}$	$\frac{\pi}{4}$	$\frac{5\pi}{16}$	$\frac{3\pi}{8}$	$\frac{7\pi}{16}$	$\frac{\pi}{2}$	$y = \cos x$	1	0.9808	0.9239	0.8315	0.7071	0.5556	0.3827	0.1951	0	06
x	0	$\frac{\pi}{16}$	$\frac{\pi}{8}$	$\frac{3\pi}{16}$	$\frac{\pi}{4}$	$\frac{5\pi}{16}$	$\frac{3\pi}{8}$	$\frac{7\pi}{16}$	$\frac{\pi}{2}$													
$y = \cos x$	1	0.9808	0.9239	0.8315	0.7071	0.5556	0.3827	0.1951	0													
c)	Attempt the following:		06																			

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228

WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code:

22201

Q. No.	Sub Q. N.	Answer	Marking Scheme												
6.	c)(i)	Using Trapezoidal rule, evaluate $\int_{-1}^1 (1+x+x^2+x^3) dx$, by taking $n = 2$.	03												
	Ans	$y = 1+x+x^2+x^3 \quad a = -1, \quad b = 1$ $\therefore h = \frac{b-a}{n} = \frac{1+1}{2} = 1$ <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>-1</td> <td>0</td> <td>1</td> </tr> <tr> <td>$y = 1+x+x^2+x^3$</td> <td>0</td> <td>1</td> <td>4</td> </tr> </table> $\int_a^b f(x) dx = \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1})]$ $a = -1, b = 1 \text{ and } h = 1$ $\therefore \int_{-1}^1 (1+x+x^2+x^3) dx = \frac{1}{2} [(0+4) + 2(1)]$ $= 3$	x	-1	0	1	$y = 1+x+x^2+x^3$	0	1	4	<p>1/2</p> <p>1</p>				
x	-1	0	1												
$y = 1+x+x^2+x^3$	0	1	4												
	ii)	Using Simpson's 1/3 rd rule evaluate $\int_1^3 \frac{dx}{x}$ taking $h = 0.5$.													
	Ans	Let $y = \frac{1}{x}, h = 0.5, a = 1, b = 3$ <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>1</td> <td>1.5</td> <td>2</td> <td>2.5</td> <td>3</td> </tr> <tr> <td>$y = \frac{1}{x}$</td> <td>1</td> <td>$\frac{2}{3}$</td> <td>$\frac{1}{2}$</td> <td>$\frac{2}{5}$</td> <td>$\frac{1}{3}$</td> </tr> </table> Using Simpson's 1/3 rd rule $\int_a^b f(x) dx = \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$ $\therefore \int_1^3 \frac{dx}{x} = \frac{0.5}{3} \left[\left(1 + \frac{1}{3}\right) + 4\left(\frac{2}{3} + \frac{2}{5}\right) + 2\left(\frac{1}{2}\right) \right]$ $\int_1^3 \frac{dx}{x} = 1.1$	x	1	1.5	2	2.5	3	$y = \frac{1}{x}$	1	$\frac{2}{3}$	$\frac{1}{2}$	$\frac{2}{5}$	$\frac{1}{3}$	<p>1/2</p> <p>1</p>
x	1	1.5	2	2.5	3										
$y = \frac{1}{x}$	1	$\frac{2}{3}$	$\frac{1}{2}$	$\frac{2}{5}$	$\frac{1}{3}$										
	<u>OR</u>	Let $y = \frac{1}{x}, h = 0.5, a = 1, b = 3$ <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>1</td> <td>1.5</td> <td>2</td> <td>2.5</td> <td>3</td> </tr> <tr> <td>$y = \frac{1}{x}$</td> <td>1</td> <td>0.6667</td> <td>0.5</td> <td>0.4</td> <td>0.3333</td> </tr> </table>	x	1	1.5	2	2.5	3	$y = \frac{1}{x}$	1	0.6667	0.5	0.4	0.3333	<p>1</p>
x	1	1.5	2	2.5	3										
$y = \frac{1}{x}$	1	0.6667	0.5	0.4	0.3333										



WINTER – 2019 EXAMINATION

Subject Name: Applied Mathematics ModelAnswer

Subject Code: 22201

Q. No.	Sub Q. N.	Answer	Marking Scheme
6.	c)(ii)	<p>Using Simpson's 1/3rd rule</p> $\int_a^b f(x) dx = \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$ $\therefore \int_1^3 \frac{dx}{x} = \frac{0.5}{3} [(1 + 0.3333) + 4(0.6667 + 0.4) + 2(0.5)]$ $\int_1^3 \frac{dx}{x} = 1.1$	<p>1</p> <p>1</p>

Important Note

In the solution of the question paper, wherever possible all the possible alternative methods of solution are given for the sake of convenience. Still student may follow a method other than the given herein. In such case, first see whether the method falls within the scope of the curriculum, and then only give appropriate marks in accordance with the scheme of marking.