



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Important Instructions to Examiners:

- 1) The answers should be examined by key words and not as word-to-word as given in the model answer scheme.
- 2) The model answer and the answer written by candidate may vary but the examiner may try to assess the understanding level of the candidate.
- 3) The language errors such as grammatical, spelling errors should not be given more Importance (Not applicable for subject English and Communication Skills).
- 4) While assessing figures, examiner may give credit for principal components indicated in the figure. The figures drawn by candidate and model answer may vary. The examiner may give credit for any equivalent figure drawn.
- 5) Credits may be given step wise for numerical problems. In some cases, the assumed constant values may vary and there may be some difference in the candidate's answers and model answer.
- 6) In case of some questions credit may be given by judgement on part of examiner of relevant answer based on candidate's understanding.
- 7) For programming language papers, credit may be given to any other program based on equivalent concept.

Q. No.	Sub Q.N.	Answers	Marking Scheme
1.		Attempt any <u>FIVE</u> of the following:	10
	a)	If $f(x) = 3x^2 - 5x + 7$ show that $f(-1) = 3f(1)$ Ans $f(-1) = 15$ $f(1) = 5$ $\therefore 3f(1) = 15$ $\therefore f(-1) = 3f(1)$	02 1 1
	b)	State whether the function $f(x) = 3x^4 + x^2 + 5 - 3\cos x + 2\sin^2 x$ is even or odd Ans $f(x) = 3x^4 + x^2 + 5 - 3\cos x + 2\sin^2 x$ $\therefore f(-x) = 3(-x)^4 + (-x)^2 + 5 - 3\cos(-x) + 2\sin^2(-x)$ $\therefore f(-x) = 3x^4 + x^2 + 5 - 3\cos x + 2\sin^2 x$ $\therefore f(-x) = f(x)$ \therefore function is an even function	02 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
c)	Find $\frac{dy}{dx}$ if $y = e^x \cdot \sin^{-1} x$ Ans $y = e^x \cdot \sin^{-1} x$ $\therefore \frac{dy}{dx} = e^x \cdot \frac{1}{\sqrt{1-x^2}} + \sin^{-1} x \cdot e^x$	02 1+1	



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
1.	c)	$\therefore \frac{dy}{dx} = e^x \left(\frac{1}{\sqrt{1-x^2}} + \sin^{-1} x \right)$	
	d)	Evaluate $e^{\int 2 \log x \, dx}$	02
	Ans	$e^{\int 2 \log x \, dx}$ $= e^{2 \int \log x \, dx}$ $= e^{2 \int \log x \cdot 1 \, dx}$ $= e^{2 \left(\log x \cdot x - \int x \cdot \frac{1}{x} \, dx \right)}$ $= e^{2 \left(x \log x - \int 1 \, dx \right)}$ $= e^{2(x \log x - x) + c}$ $= e^{2x(\log x - 1) + c}$	<p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>
e)	Evaluate $\int \sin^2 x \, dx$	02	
Ans	$\int \sin^2 x \, dx$ $= \frac{1}{2} \int 2 \sin^2 x \, dx$ $= \frac{1}{2} \int (1 - \cos 2x) \, dx$ $= \frac{1}{2} \left(x - \frac{\sin 2x}{2} \right) + c$	<p>1</p> <p>1</p>	
f)	Find the area under the curve $y = x^2$ from $x = 0$ to $x = 3$ with x -axis.	02	
Ans	$\text{Area } A = \int_a^b y \, dx$ $= \int_0^3 x^2 \, dx$ $= \left[\frac{x^3}{3} \right]_0^3$	<p>1/2</p> <p>1/2</p>	

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
1.	f)	$= \left(\frac{3^3}{3} - 0 \right)$ $= 9$	½
	g)	Express $z = 1 - i$ in Polar form.	02
	Ans	$z = 1 - i$ $\therefore x = 1, y = -1$ $\therefore r = z = \sqrt{x^2 + y^2} = \sqrt{(1)^2 + (-1)^2} = \sqrt{2}$ $\theta = 2\pi - \tan^{-1} \left(\left \frac{y}{x} \right \right) = 2\pi - \tan^{-1} \left(\left \frac{-1}{1} \right \right) = 2\pi - \tan^{-1}(1)$ $\theta = 2\pi - \frac{\pi}{4} = \frac{7\pi}{4}$ $\therefore \text{Polar form is } z = r(\cos \theta + i \sin \theta)$ $\therefore 1 - i = \sqrt{2} \left(\cos \left(\frac{7\pi}{4} \right) + i \sin \left(\frac{7\pi}{4} \right) \right)$	½
2.		Attempt any <u>THREE</u> of the following:	12
	a)	Find $\frac{dy}{dx}$ if $x^2 + y^2 = 4xy$	04
	Ans	$x^2 + y^2 = 4xy$ $\therefore 2x + 2y \frac{dy}{dx} = 4 \left(x \frac{dy}{dx} + y \cdot 1 \right)$ $\therefore 2x + 2y \frac{dy}{dx} = 4x \frac{dy}{dx} + 4y$ $\therefore (2y - 4x) \frac{dy}{dx} = 4y - 2x$ $\therefore \frac{dy}{dx} = \frac{4y - 2x}{2y - 4x}$ $\therefore \frac{dy}{dx} = \frac{2y - x}{y - 2x}$	2
			1
			1



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q. N.	Answers	Marking Scheme
2.	b)	<p>If $x = a(\theta + \sin \theta)$, $y = a(1 - \cos \theta)$, find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{2}$</p> <p>Ans $x = a(\theta + \sin \theta)$ $y = a(1 - \cos \theta)$</p> $\frac{dx}{d\theta} = a(1 + \cos \theta)$ $\frac{dy}{d\theta} = a(0 + \sin \theta) = a \sin \theta$ $\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{a \sin \theta}{a(1 + \cos \theta)}$ $\frac{dy}{dx} = \frac{\sin \theta}{1 + \cos \theta}$ <p>at $\theta = \frac{\pi}{2}$</p> $\therefore \frac{dy}{dx} = \frac{\sin \frac{\pi}{2}}{1 + \cos \frac{\pi}{2}} = 1$	<p>04</p> <p>1+1</p> <p>1</p> <p>1</p>
	c)	<p>Find the radius of curvature of the curve $\sqrt{x} + \sqrt{y} = 1$ at $\left(\frac{1}{4}, \frac{1}{4}\right)$</p> <p>Ans $\sqrt{x} + \sqrt{y} = 1$</p> $\therefore \frac{1}{2\sqrt{x}} + \frac{1}{2\sqrt{y}} \frac{dy}{dx} = 0$ $\therefore \frac{dy}{dx} = -\frac{\sqrt{y}}{\sqrt{x}}$ $\therefore \frac{d^2y}{dx^2} = -\frac{\left(\sqrt{x} \frac{1}{2\sqrt{y}} \frac{dy}{dx} - \sqrt{y} \frac{1}{2\sqrt{x}}\right)}{x}$ $\therefore \frac{d^2y}{dx^2} = -\frac{\left(\frac{\sqrt{x}}{2\sqrt{y}} \cdot -\frac{\sqrt{y}}{\sqrt{x}} - \frac{\sqrt{y}}{2\sqrt{x}}\right)}{x} = -\frac{\left(\frac{-1}{2} - \frac{\sqrt{y}}{2\sqrt{x}}\right)}{x}$ <p>at $\left(\frac{1}{4}, \frac{1}{4}\right)$</p>	<p>04</p> <p>½</p> <p>½</p> <p>½</p> <p>½</p>

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
2.	d)	$\therefore \frac{dy}{dx} = -\frac{\sqrt{1/4}}{\sqrt{1/4}} = -1$ $\therefore \frac{d^2y}{dx^2} = -\frac{\left(\frac{-1}{2} - \frac{\sqrt{1/4}}{2\sqrt{1/4}}\right)}{1/4} = 4$ $\therefore \text{Radius of curvature} = \frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{3/2}}{\frac{d^2y}{dx^2}} = \frac{\left[1 + (-1)^2\right]^{3/2}}{4}$ $= 0.707$	<p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>
		<p>Find the maximum and minimum value of $x^3 - 9x^2 + 24x$</p> <p>Let $y = x^3 - 9x^2 + 24x$</p> <p>d) $\therefore \frac{dy}{dx} = 3x^2 - 18x + 24$</p> <p>Ans $\therefore \frac{d^2y}{dx^2} = 6x - 18$</p> <p>Consider $\frac{dy}{dx} = 0$</p> $3x^2 - 18x + 24 = 0$ <p>$\therefore x = 2$ or $x = 4$</p> <p>at $x = 2 \quad \therefore \frac{d^2y}{dx^2} = 6(2) - 18 = -6 < 0$</p> <p>$\therefore y$ is maximum at $x = 2$</p> $y_{\max} = (2)^3 - 9(2)^2 + 24(2) = 20$ <p>at $x = 4 \quad \therefore \frac{d^2y}{dx^2} = 6(4) - 18 = 6 > 0$</p> <p>$\therefore y$ is minimum at $x = 4$</p> $y_{\min} = (4)^3 - 9(4)^2 + 24(4) = 16$	<p>04</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>
		<p>Note: If students attempted to solve the question give appropriate marks.</p>	

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
3.	b)	Find $\frac{dy}{dx}$ if $y = x^x + (\sin x)^x$	04
	Ans	<p>Let $u = x^x$</p> <p>Taking log on both sides,</p> $\therefore \log u = \log x^x$ $\therefore \log u = x \log x$ $\therefore \frac{1}{u} \frac{du}{dx} = x \frac{1}{x} + \log x \cdot 1$ $\therefore \frac{1}{u} \frac{du}{dx} = 1 + \log x$ $\therefore \frac{du}{dx} = u(1 + \log x)$ <p>Let $v = (\sin x)^x$</p> <p>taking log on both sides,</p> $\therefore \log v = \log (\sin x)^x$ $\therefore \log v = x \log \sin x$ $\therefore \frac{1}{v} \frac{dv}{dx} = x \frac{1}{\sin x} \cdot \cos x + \log \sin x \cdot 1$ $\therefore \frac{1}{v} \frac{dv}{dx} = x \cot x + \log \sin x$ $\therefore \frac{dv}{dx} = v(x \cot x + \log \sin x)$ $\therefore \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$ $\therefore \frac{dy}{dx} = x^x (1 + \log x) + (\sin x)^x (x \cot x + \log \sin x)$	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p>
	c)	If $y = e^{3\sec x + 4\tan x}$ find $\frac{dy}{dx}$	04
	Ans	<p>$y = e^{3\sec x + 4\tan x}$</p> $\therefore \frac{dy}{dx} = e^{3\sec x + 4\tan x} (3\sec x \cdot \tan x + 4\sec^2 x)$	4



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
3.	d)	Evaluate $\int \frac{\sec^2 x}{(1 + \tan x)(3 + \tan x)} dx$	04
	Ans	$\int \frac{\sec^2 x}{(1 + \tan x)(3 + \tan x)} dx$ <p>Put $\tan x = t$</p> $\therefore \sec^2 x dx = dt$ $\therefore \int \frac{1}{(1+t)(3+t)} dt$ $\frac{1}{(1+t)(3+t)} = \frac{A}{1+t} + \frac{B}{3+t}$ $\therefore 1 = A(3+t) + B(1+t)$ $\therefore \text{Put } t = -1, A = \frac{1}{2}$ $\text{Put } t = -3, B = \frac{-1}{2}$ $\therefore \frac{1}{(1+t)(3+t)} = \frac{1/2}{1+t} - \frac{1/2}{3+t}$ $\therefore \int \frac{1}{(1+t)(3+t)} dt = \int \left(\frac{1/2}{1+t} - \frac{1/2}{3+t} \right) dt$ $= \frac{1}{2} \log(1+t) - \frac{1}{2} \log(3+t) + c$ $= \frac{1}{2} \log(1 + \tan x) - \frac{1}{2} \log(3 + \tan x) + c$ <p>OR</p> $\int \frac{\sec^2 x}{(1 + \tan x)(3 + \tan x)} dx$ <p>Put $\tan x = t$</p> $\therefore \sec^2 x dx = dt$ $\int \frac{1}{(1+t)(3+t)} dt$	<p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p>

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
3.	d)	$= \int \frac{1}{t^2 + 4t + 3} dt$ $\text{Third Term} = \frac{4^2}{4} = 4$ $= \int \frac{1}{t^2 + 4t + 4 - 4 + 3} dt$ $= \int \frac{1}{(t+2)^2 - 1} dt$ $= \frac{1}{2} \log \left \frac{t+2-1}{t+2+1} \right + c$ $= \frac{1}{2} \log \left \frac{t+1}{t+3} \right + c$ $= \frac{1}{2} \log \left \frac{\tan x + 1}{\tan x + 3} \right + c$	<p>1/2</p> <p>1</p> <p>1</p> <p>1/2</p>
4.		<p>Attempt any <u>THREE</u> of the following:</p> <p>a) Evaluate $\int x \tan^{-1} x dx$</p> <p>Ans $\int \tan^{-1} x \cdot x dx$</p> $= \tan^{-1} x \int x dx - \int \left(\int x dx \frac{d}{dx} (\tan^{-1} x) \right) dx$ $= \tan^{-1} x \frac{x^2}{2} - \int \frac{x^2}{2} \frac{1}{1+x^2} dx$ $= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} \int \frac{x^2}{1+x^2} dx$ $= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} \int \left(\frac{1+x^2-1}{1+x^2} \right) dx$ $= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} \int \left(\frac{1+x^2}{1+x^2} - \frac{1}{1+x^2} \right) dx$ $= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} \int \left(1 - \frac{1}{1+x^2} \right) dx$ $= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} (x - \tan^{-1} x) + c$	<p>12</p> <p>04</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p>

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
4.	b)	<p>Evaluate $\int \frac{dx}{4+5\cos x}$</p> <p>Ans $\int \frac{dx}{4+5\cos x}$</p> <p>Put $\tan \frac{x}{2} = t$, $dx = \frac{2dt}{1+t^2}$, $\cos x = \frac{1-t^2}{1+t^2}$</p> $= \int \frac{\frac{2dt}{1+t^2}}{4+5\left(\frac{1-t^2}{1+t^2}\right)}$ $= \int \frac{2dt}{4(1+t^2)+5(1-t^2)}$ $= 2 \int \frac{dt}{4+4t^2+5-5t^2}$ $= 2 \int \frac{dt}{9-t^2}$ $= 2 \int \frac{dt}{(3)^2-t^2}$ $= 2 \frac{1}{2(3)} \log \left \frac{3+t}{3-t} \right + c$ $= \frac{1}{3} \log \left \frac{3+\tan \frac{x}{2}}{3-\tan \frac{x}{2}} \right + c$	<p>04</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p>
	c)	<p>Evaluate $\int \frac{2x^2+5}{(x-1)(x+2)(x+3)} dx$</p> <p>Ans $\int \frac{2x^2+5}{(x-1)(x+2)(x+3)} dx$</p> <p>Consider $\frac{2x^2+5}{(x-1)(x+2)(x+3)} = \frac{A}{x-1} + \frac{B}{x+2} + \frac{C}{x+3}$</p> $\therefore 2x^2+5 = A(x+2)(x+3) + B(x-1)(x+3) + C(x-1)(x+2)$ <p>Put $x = 1$</p> $\therefore 7 = 12A$	<p>04</p> <p>1/2</p>

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
4.	c)	$\therefore A = \frac{7}{12}$ $\text{Put } x = -2 \quad \Rightarrow 13 = -3B$ $\therefore B = \frac{-13}{3}$ $\text{Put } x = -3 \quad \Rightarrow 23 = 4C$ $\therefore C = \frac{23}{4}$ $\therefore \int \frac{2x^2 + 5}{(x-1)(x+2)(x+3)} dx = \int \left(\frac{\frac{7}{12}}{x-1} + \frac{\frac{-13}{3}}{x+2} + \frac{\frac{23}{4}}{x+3} \right) dx$ $= \frac{7}{12} \log(x-1) - \frac{13}{3} \log(x+2) + \frac{23}{4} \log(x+3) + c$	<p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2+1/2+1/2</p>
	d)	<p>Evaluate $\int \frac{1}{\sqrt{16-6x-x^2}} dx$</p> <p>Ans $\int \frac{1}{\sqrt{16-6x-x^2}} dx$</p> <p>Third Term = $\frac{(6)^2}{4} = 9$</p> $= \int \frac{1}{\sqrt{16+9-9-6x-x^2}} dx$ $= \int \frac{1}{\sqrt{25-(9+6x+x^2)}} dx$ $= \int \frac{1}{\sqrt{(5)^2-(x+3)^2}} dx$ $= \sin^{-1}\left(\frac{x+3}{5}\right) + c$ <p>OR</p> $\int \frac{1}{\sqrt{16-6x-x^2}} dx$	<p>04</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
4.	d)	$= \int \frac{1}{\sqrt{-(x^2 + 6x - 16)}} dx$ $\text{Third Term} = \frac{(6)^2}{4} = 9$ $= \int \frac{1}{\sqrt{-(x^2 + 6x + 9 - 9 - 16)}} dx$ $= \int \frac{1}{\sqrt{-(x^2 + 6x + 9 - 25)}} dx$ $= \int \frac{1}{\sqrt{-(x+3)^2 - (5)^2}} dx$ $= \int \frac{1}{\sqrt{(5)^2 - (x+3)^2}} dx$ $= \sin^{-1}\left(\frac{x+3}{5}\right) + c$	<p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1</p>
	e)	<p>Evaluate $\int_0^{\pi/2} \frac{dx}{1 + \cot x}$</p> <p>Ans $\int_0^{\pi/2} \frac{dx}{1 + \cot x}$</p> $\therefore I = \int_0^{\pi/2} \frac{1}{1 + \frac{\cos x}{\sin x}} dx$ $\therefore I = \int_0^{\pi/2} \frac{\sin x}{\sin x + \cos x} dx \text{ -----(1)}$ $\therefore I = \int_0^{\pi/2} \frac{\sin(\pi/2 - x)}{\sin(\pi/2 - x) + \cos(\pi/2 - x)} dx$ $\therefore I = \int_0^{\pi/2} \frac{\cos x}{\cos x + \sin x} dx \text{ -----(2)}$ <p>add (1) and (2)</p>	<p>04</p> <p>1/2</p> <p>1</p>

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
4.	e)	$\therefore I + I = \int_0^{\pi/2} \frac{\sin x}{\sin x + \cos x} dx + \int_0^{\pi/2} \frac{\cos x}{\cos x + \sin x} dx$ $2I = \int_0^{\pi/2} \frac{\sin x + \cos x}{\sin x + \cos x} dx$ $2I = \int_0^{\pi/2} 1 dx$ $2I = [x]_0^{\pi/2}$ $2I = \frac{\pi}{2} - 0$ $\therefore I = \frac{\pi}{4}$	<p>1/2</p> <p>1</p> <p>1</p>
5.		<p>Attempt any TWO of the following:</p> <p>a) Find the area between the curves $y = x$ and $y = x^2$</p> <p>Ans</p> $y = x$ $y = x^2$ $\therefore x - x^2 = 0$ $\therefore x(1 - x) = 0$ $\therefore x = 0, 1$ $\therefore A = \int_0^1 (x - x^2)$ $A = \left(\frac{x^2}{2} - \frac{x^3}{3} \right)_0^1$ $A = \left(\frac{1}{2} - \frac{1}{3} \right)$ $\therefore A = \frac{1}{6} \text{ or } 0.167$	<p>04</p> <p>2</p> <p>1</p> <p>1</p> <p>1</p>
5.		<p>Attempt the following</p> <p>b)(i) Find the order and degree of the differential equation</p>	<p>12</p>

OUR CENTERS :



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
5.	b)	$\frac{d^2y}{dx^2} = \sqrt{1 + \frac{dy}{dx}}$	03
	(i)	$\frac{d^2y}{dx^2} = \sqrt{1 + \frac{dy}{dx}}$	
	Ans	Squaring both sides, we get $\left(\frac{d^2y}{dx^2}\right)^2 = 1 + \frac{dy}{dx}$ $\therefore \text{Order} = 2$ $\text{Degree} = 2$	
	b)(ii)	Solve $\frac{dy}{dx} + y \cot x = \cos ecx$ Ans $\frac{dy}{dx} + y \cot x = \cos ecx$ Comparing with $\frac{dy}{dx} + Py = Q$ $P = \cot x$, $Q = \cos ecx$ Integrating factor $IF = e^{\int \cot x dx} = e^{\log(\sin x)} = \sin x$ $y \cdot IF = \int Q \cdot IF dx + c$ $\therefore y \sin x = \int \cos ecx \cdot \sin x dx$ $\therefore y \sin x = \int 1 dx$ $\therefore y \sin x = x + c$	03
	c)	If $L \frac{di}{dt} = 30 \cdot \sin(10\pi t)$, find i in terms of t , given that $L = 2$ and $i = 0$ at $t = 0$	06
	Ans	$L di = 30 \sin(10\pi t) dt$ $\int L di = \int 30 \sin(10\pi t) dt$ $Li = 30 \left(\frac{-\cos(10\pi t)}{10\pi} \right) + c$	

OUR CENTERS :



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
5.	c)	$Li = \frac{-3\cos(10\pi t)}{\pi} + c$ <p>at $t = 0, i = 0$</p> $L(0) = \frac{-3\cos(0)}{\pi} + c$ $0 = \frac{-3}{\pi} + c$ $\therefore c = \frac{3}{\pi}$ $\therefore Li = \frac{-3\cos(10\pi t)}{\pi} + \frac{3}{\pi}$ <p>at $L = 2$</p> $\therefore 2i = \frac{-3\cos(10\pi t)}{\pi} + \frac{3}{\pi}$ $\therefore i = \frac{3}{2\pi}(-\cos(10\pi t) + 1)$	<p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p>
6.		<p>Attempt any <u>TWO</u> of the following:</p> <p>a) Attempt the following</p> <p>(i) Express $\frac{2-\sqrt{3}i}{1+i}$ in $x+iy$ form.</p> <p>Ans</p> $\frac{2-\sqrt{3}i}{1+i}$ $= \frac{2-\sqrt{3}i}{1+i} \times \frac{1-i}{1-i}$ $= \frac{2-2i-\sqrt{3}i+\sqrt{3}i^2}{1-i^2}$ $= \frac{2-(2+\sqrt{3})i+\sqrt{3}(-1)}{1-i^2}$ $= \frac{2-(2+\sqrt{3})i-\sqrt{3}}{1+1}$	<p>04</p> <p>03</p> <p>1</p> <p>1/2</p> <p>1/2</p>



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22210

Q. No.	Sub Q.N.	Answers	Marking Scheme
6.	a)(i)	$\frac{(2-\sqrt{3})-(2+\sqrt{3})i}{2}$ $= \frac{(2-\sqrt{3})}{2} - \frac{(2+\sqrt{3})i}{2}$	1
	a)(ii)	<p>Find $L\{e^{-4t}t^2\}$</p> <p>Ans $L\{e^{-4t}t^2\}$</p> $L\{t^2\} = \frac{2!}{s^3} = \frac{2}{s^3}$ $\therefore L\{e^{-4t}t^2\} = \frac{2}{(s+4)^3}$	03
	b)	<p>Find $L^{-1}\left\{\frac{2s^2-4}{(s+1)(s-2)(s-3)}\right\}$</p> <p>Ans $L^{-1}\left\{\frac{2s^2-4}{(s+1)(s-2)(s-3)}\right\}$</p> <p>Let</p> $\frac{2s^2-4}{(s+1)(s-2)(s-3)} = \frac{A}{s+1} + \frac{B}{s-2} + \frac{C}{s-3}$ $2s^2-4 = (s-2)(s-3)A + (s+1)(s-3)B + (s+1)(s-2)C$ <p>Put $s = -1$</p> $\therefore -2 = 12A$ $\therefore A = -\frac{1}{6}$ <p>Put $s = 2$</p> $4 = -3B$ $\therefore B = -\frac{4}{3}$ <p>Put $s = 3$</p> $14 = 4C$	04

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228

