

APPLIED MATHEMATICS SOLVED PAPER – MAY 2018

N.B:- (1) Question no. 1 is compulsory.
(2) Attempt any 3 questions from remaining five questions.

Q.1.(a) Evaluate $\int_0^{\infty} 5^{-4x^2} dx$ [3]

Ans: Let $I = \int_0^{\infty} 5^{-4x^2} dx$

put $5^{-4x^2} = e^{-t}$

taking log on both sides,

$$4x^2 \log 5 = t$$

$$x^2 = \frac{t}{4 \log 5} \Rightarrow x = \frac{\sqrt{t}}{2\sqrt{\log 5}}$$

diff. w.r.t x,

$$dx = \frac{t^{-1/2}}{4\sqrt{\log 5}} dt \quad \text{lim} \rightarrow [0, \infty]$$

$$\therefore I = \int_0^{\infty} \frac{e^{-t}}{4\sqrt{\log 5}} t^{-1/2}$$

$$\therefore I = \frac{1}{4\sqrt{\log 5}} \int_0^{\infty} e^{-t} \cdot t^{-1/2} dt$$

$$\boxed{\therefore I = \frac{\sqrt{\pi}}{4\sqrt{\log 5}}}$$

$$\dots\dots\dots\{ \int_0^{\infty} e^{-t} \cdot t^{-1/2} dt = \sqrt{\pi} \}$$

(b) Solve $\frac{dy}{dx} = x \cdot y$ with help of Euler's method ,given that $y(0)=1$ and find

y when $x=0.3$ [3]

(Take $h=0.1$)

Ans : $\frac{dy}{dx} = x \cdot y = f(x, y) \quad x_0 = 0, y_0 = 1$

$$y_n = y_{n-1} + h \cdot f(x_{n-1}, y_{n-1})$$

OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228

Iteration (n)	x_n	y_n	$f(x_n, y_n)$	y_{n+1} $= y_n$ $+ h \cdot f(x_n, y_n)$
0	0	1	0	1
1	0.1	1	0.1	1.01
2	0.2	1.01	0.202	1.0302

$$\therefore y(0.3) = 1.0302$$

(c) Evaluate $\frac{d^4 y}{dx^4} + 2\frac{d^2 y}{dx^2} + y = 0$ [3]

Ans: put $\frac{d}{dx} = D$

$$\therefore D^4 y + 2D^2 y + y = 0$$

$$\therefore D^4 + 2D^2 + 1 = 0$$

Put $D^2 = t$

$$\Rightarrow t^2 + 2t + 1 = 0$$

$$\Rightarrow t = -1, -1$$

Roots are : $D = +i, -i, +i, -i$

The complementary solution of given eqn is

$$y_c = y_g = (C_1 + xC_2)\cos x + (C_3 + xC_4)\sin x$$

(d) Evaluate $\int_0^1 \sqrt{\sqrt{x} - x} dx$ [3]

Ans: Let $I = \int_0^1 \sqrt{\sqrt{x} - x} dx$

$$I = \int_0^1 \sqrt{(\sqrt{x} - \sqrt{x}) \cdot \sqrt{x}} dx$$

Take \sqrt{x} common,

$$I = \int_0^1 x^{1/4} \sqrt{1 - x^{1/2}} dx$$

Put $x^{1/2} = t$

Squaring both sides,

$$\therefore x = t^2$$

Differentiate w.r.t x,

$$\therefore dx = 2t \cdot dt$$

Limits after substitution : Lim $\rightarrow [0, 1]$

$$\therefore I = \int_0^1 t^{1/2} \sqrt{1-t} \cdot 2 \cdot t \, dt$$

$$= 2 \int_0^1 t^{3/2} \sqrt{1-t} \, dt$$

$$= 2 \beta\left(\frac{5}{2}, \frac{3}{2}\right)$$

$$\dots\dots\left\{ \int_0^1 t^m \cdot (1-t)^n = \beta(m+1, n+1) \right\}$$

$$\boxed{\therefore I = \frac{\pi}{8}}$$

(e) Solve: $(1 + \log x \cdot y) dx + (1 + \frac{x}{y}) dy = 0$

[4]

Ans : Compare given eqn with $Mdx + Ndy = 0$

$$\therefore M = (1 + \log x \cdot y) \quad \therefore N = 1 + \frac{x}{y}$$

$$\frac{\partial M}{\partial y} = \frac{1}{xy} \cdot x = \frac{1}{y}$$

$$\frac{\partial N}{\partial x} = \frac{1}{y}$$

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Hence the given differential eqn is exact.

The solution of exact differential eqn is given by,

$$\int M dx + \int \left(N - \frac{\partial}{\partial y} \int M dx \right) dy = c \quad \text{-----(1)}$$

$$\int M dx = \int (1 + \log xy) dx = x + \log xy \cdot x - x = x \cdot \log xy$$

$$\frac{\partial}{\partial y} \int M dx = \frac{x}{y}$$

$$\int \left(N - \frac{\partial}{\partial y} \int M dx \right) dy = \int \left(1 + \frac{x}{y} - \frac{x}{y} \right) dy = y$$

From eqn (1), the solution of given differential eqn is ,

$$\boxed{x \cdot \log xy + y = c}$$

(f) Evaluate $I = \int_0^1 \int_0^{\sqrt{1+x^2}} \frac{dx \cdot dy}{1+x^2+y^2}$

[4]

Ans :

$$I = \int_0^1 \int_0^{\sqrt{1+x^2}} \frac{dx \cdot dy}{1+x^2+y^2}$$

$$I = \int_0^1 \frac{1}{\sqrt{1+x^2}} \left[\tan^{-1} \frac{y}{\sqrt{1+x^2}} \right]_0^{\sqrt{1+x^2}} dx$$

$$\therefore I = \int_0^1 \frac{\pi}{4} \frac{1}{\sqrt{1+x^2}} dx$$

$$\therefore I = \frac{\pi}{4} \left[\log (x + \sqrt{1+x^2}) \right]_0^1$$

$$\therefore I = \frac{\pi}{4} \log(1 + \sqrt{2})$$

Q.2. (a) Solve $xy(1+xy^2)\frac{dy}{dx} = 1$

[6]

Ans:

$$\therefore \frac{dx}{dy} = xy + x^2y^3$$

$$\therefore \frac{1}{x^2} \frac{dx}{dy} - \frac{1}{x}y = y^3$$

$$\therefore \frac{dv}{dy} + vy = y^3$$

Now, put $-\frac{1}{x} = v$

..... $\left(\frac{1}{x^2} \frac{dx}{dy} = \frac{dv}{dy} \right)$

This is linear differential eqn.

\therefore Integrating Factor = $e^{\int y dy} = e^{\frac{y^2}{2}}$

The solution of linear diff. eqn is given by,

$$v \cdot (I.F.) = \int (I.F.) (R.H.S) + c$$

$$v e^{\frac{y^2}{2}} = \int e^{\frac{y^2}{2}} \cdot y^3 dy + c$$

$$-\frac{1}{x} e^{\frac{y^2}{2}} = e^{\frac{y^2}{2}} (y^2 - 2) + c$$


Where c is constant of integration.

(b) Find the area inside the circle $r=a \sin\theta$ and outside the cardioide $r=a(1+\cos\theta)$

[6]

Ans : Intersection of cardioide and circle is,

$$r=a(1+\cos\theta) \text{ and } r=asin\theta$$



$$a \sin \theta = a(1 + \cos \theta) \Rightarrow \theta = 90^\circ$$

$$a(1 + \cos \theta) \leq r \leq a \sin \theta$$

$$\frac{\pi}{2} \leq \theta \leq \pi$$

Area of region bounded by given circle and cardioid ,

$$\begin{aligned} I &= \int_{\frac{\pi}{2}}^{\pi} \int_{a \sin \theta}^{a(1 + \cos \theta)} r \, dr \, d\theta \\ &= \int_{\frac{\pi}{2}}^{\pi} \frac{a^2}{2} (\sin^2 \theta - 1 - 2 \cos \theta - \cos^2 \theta) \, d\theta \\ &= \int_{\frac{\pi}{2}}^{\pi} \frac{a^2}{2} (-1 - 2 \cos \theta - \cos 2\theta) \, d\theta \\ &= \frac{a^2}{2} \left[-\theta - 2 \sin \theta - \frac{\sin 2\theta}{2} \right]_{\frac{\pi}{2}}^{\pi} \\ I &= \frac{a^2}{2} \left[(-\pi - 0 - 0) - \left(-\frac{\pi}{2} - 2 - 0\right) \right] \end{aligned}$$

Required area is $I = \frac{a^2}{2} \left(2 - \frac{\pi}{2} \right)$

(c) Apply Rungee-Kutta Method of fourth order to find an approximate value of y when x=0.2 given that $\frac{dy}{dx} = x + y$ when y=1 at x=0 with step size h=0.2. [8]

Ans: $\frac{dy}{dx} = x + y \quad x_0 = 0, \quad y_0 = 1, \quad h = 0.2$

$$f(x, y) = x + y$$

$$k_1 = h \cdot f(x_0, y_0) = 0.2 f(0, 1) = 0.2$$

$$k_2 = h \cdot f\left(x_0 + \frac{h}{2}, y_0 + \frac{k_1}{2}\right) = 0.2 f(0.1, 1.1) = 0.24$$

$$k_3 = h.f(x_0 + \frac{h}{2}, y_0 + \frac{k_2}{2}) = 0.2f(0.1, 1.12) = 0.244$$

$$k_4 = h.f(x_0 + h, y_0 + k_3) = 0.2 f(0.2, 1.244) = 0.2888$$

$$k = \frac{k_1+2k_2+2k_3+k_4}{6} = \frac{0.24+0.48+0.488+0.2888}{6} = 0.2428$$

The value of y at x=0.2 is given by,


$$y(0.2) = y_0 + k = 1 + 0.2428$$

$$y(0.2) = 1.2428$$

Q.3 (a) Show that the length of curve $9ay^2=x(x-3a)^2$ is $4\sqrt{3}a$. [6]

Ans : Curve : $9ay^2=x(x-3a)^2$ (1)

The given curve is strophoid.



Differentiate eqn (1) w.r.t x,

$$18ay \frac{dy}{dx} = 2x(x-3a) + (x-3a)^2$$

$$\therefore 18ay \frac{dy}{dx} = 3(x-3a)(x-a)$$

$$\therefore \frac{dy}{dx} = \frac{(x-3a)(x-a)}{6ay}$$

Squaring both the sides,

$$\left(\frac{dy}{dx}\right)^2 = \frac{(x-3a)^2(x-a)^2}{36 a^2 y^2}$$

$$\therefore \left(\frac{dy}{dx}\right)^2 = \frac{(x-3a)^2(x-a)^2}{4a x (x-3a)^2} \quad \text{..... from (1)}$$

$$\therefore \left(\frac{dy}{dx}\right)^2 = \frac{(x-a)^2}{4ax}$$

The perimeter of given curve is ,

$$S = \int_0^{3a} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_0^{3a} \sqrt{1 + \frac{(x-a)^2}{4ax}} dx = \int_0^{3a} \sqrt{\frac{(x+a)^2}{4ax}} dx$$

$$\therefore S = \int_0^{3a} \frac{x+a}{2\sqrt{x}\sqrt{a}} dx$$

$$\therefore S = \frac{1}{2\sqrt{a}} \int_0^{3a} \frac{x+a}{\sqrt{x}} dx$$

$$= \frac{1}{2\sqrt{a}} \left[\frac{2x\sqrt{x}}{3} + 2\sqrt{x} \right]_0^{3a}$$

$$= \frac{1}{2\sqrt{a}} \left(\frac{2a\sqrt{3a}}{1} + 2\sqrt{3a} \right)$$

$$\therefore S = 2\sqrt{3} \quad \text{----- (Half curve length)}$$

∴ The total length of given curve = 2 S = 4 √3 units.

(b) Change the order of integration of $\int_0^1 \int_{-\sqrt{2y-y^2}}^{1+\sqrt{1-y^2}} f(x,y) dx dy$. [6]

Ans: Let $I = \int_0^1 \int_{-\sqrt{2y-y^2}}^{1+\sqrt{1-y^2}} f(x,y) dx dy$

Region of integration : $-\sqrt{2y-y^2} \leq x \leq 1 + \sqrt{1-y^2}$
 $0 \leq y \leq 1$

Curves : (i) $x = -\sqrt{2y-y^2} \Rightarrow x^2 + y^2 = 2y \Rightarrow x^2 + (y-1)^2 = 1$


Circle with centre (0,1) and radius 1.

(ii) $x = 1 + \sqrt{1-y^2} \Rightarrow (x-1)^2 + y^2 = 1$

Circle with centre (1,0) and radius 1.

(iii) $y = 0$ line i.e equation of x - axis.

(iv) $y = 1$ line parallel to x - axis.



OUR CENTERS :

KALYAN | DOMBIVLI | THANE | NERUL | DADAR

Contact - 9136008228

Divide the region R into R1 and R2

$$\therefore R = R1 \cup R2$$

After changing the order of integration ,

$$\text{For region R1 : } 0 \leq y \leq 1 - \sqrt{1 - x^2}$$

$$0 \leq x \leq 1$$

$$\text{For region R2 : } 0 \leq y \leq \sqrt{1 - (x - 1)^2}$$

$$1 \leq x \leq 2$$


As the region is divided in two parts the integration will be the union of the two region limits.

$$I = \int_0^1 \int_0^{1-\sqrt{1-x^2}} f(x,y) dy dx + \int_1^2 \int_0^{\sqrt{1-(x-1)^2}} f(x,y) dy dx$$

This is the integration after changing order from dx dy to dy dx of given integration region.

(c) Find the volume of the paraboloid $x^2 + y^2 = 4z$ cut off by the plane $z = 4$ [8]

Ans: Paraboloid : $x^2 + y^2 = 4z$ Plane : $z = 4$



Cartesian coordinate \longrightarrow cylindrical coordinates

$$(x, y, z) \longrightarrow (r, \theta, z)$$

Put $x = r \cos \theta$, $y = r \sin \theta$, $z = z$ $\therefore x^2 + y^2 = r^2$

\therefore Paraboloid : $r^2 = 4z$ and Plane : $z = 4$

If we are passing one arrow parallel to z axis from -ve to +ve we will get limits of z

$$\therefore \frac{r^2}{4} \leq z \leq 4$$

$$0 \leq r \leq 4$$

$$0 \leq \theta \leq \frac{\pi}{2}$$

Volume of given paraboloid cut off by the plane is given by ,

$$\begin{aligned} V &= 4 \int_0^{\frac{\pi}{2}} \int_0^4 \int_{\frac{r^2}{4}}^4 r \, dr \, d\theta \, dz \\ &= 4 \int_0^{\frac{\pi}{2}} \int_0^4 \left[4r - \frac{r^4}{16} \right] \frac{4}{4} \, dr \, d\theta \\ &= 4 \int_0^{\frac{\pi}{2}} \int_0^4 \left(4r - \frac{r^3}{4} \right) \, dr \, d\theta \\ &= 4 \int_0^{\frac{\pi}{2}} \left[2r^2 - \frac{r^4}{16} \right]_0^4 \, d\theta \\ &= 4 \int_0^{\frac{\pi}{2}} (32 - 16) \, d\theta \end{aligned}$$

$V = 32 \pi$ cubic units

Q.4 (a) Show that $\int_0^1 \frac{x^a - 1}{\log x} \, dx = \log(a+1)$ [6]

Ans : let $I = \int_0^1 \frac{x^a - 1}{\log x} \, dx$

Taking 'a' as parameter ,

$$I(a) = \int_0^1 \frac{x^a - 1}{\log x} \, dx \quad \text{----- (1)}$$

differentiate w.r.t a ,

$$\frac{dI(a)}{da} = \frac{d}{da} \int_0^1 \frac{x^a - 1}{\log x} \, dx$$

$$\therefore \frac{dI(a)}{da} = \int_0^1 \frac{\partial}{\partial a} \frac{x^a - 1}{\log x} \, dx \quad \text{.....\{ D.U.I.S f(x)\}}$$

$$\therefore \frac{dI(a)}{da} = \int_0^1 \frac{x^a \cdot \log x}{\log x} dx \quad \dots\dots\dots \left\{ \frac{dx^a}{da} = x^a \cdot \log a \right\}$$

$$\therefore \frac{dI(a)}{da} = \int_0^1 x^a dx$$

$$\therefore \frac{dI(a)}{da} = \left[\frac{x^{a+1}}{a+1} \right]_0^1$$

$$\therefore \frac{dI(a)}{da} = \frac{1}{a+1} - 0$$

$$\therefore \frac{dI(a)}{da} = \frac{1}{a+1}$$

now , integrate w.r.t a,

$$I(a) = \int \frac{1}{a+1} da$$

$$I(a) = \log(a+1) + c \quad \text{----- (2)}$$

where c is constant of integration

put a=0 in eqn (1),

$$I(0) = \int_0^1 0 dx = 0$$

And

From eqn (2), $I(0) = c$

$$\therefore c = 0$$

$$\therefore I = \log(a+1)$$

Hence proved.

(b) If y satisfies the equation $\frac{dy}{dx} = x^2y - 1$ with $x_0 = 0, y_0 = 1$ using Taylor's Series Method find y at $x = 0.1$ (take $h=0.1$). [6]

Ans : $\frac{dy}{dx} = x^2y - 1 \quad x_0 = 0, y_0 = 1, h = 0.1$

To find : $y(0.1)$

$$y' = x^2y - 1 \quad , \quad y'_0 = -1$$

$$y'' = x^2y' + 2xy \quad , \quad y''_0 = 0$$

$$y''' = x^2y'' + 2y'x + 2y + 2xy' \quad , \quad y'''_0 = 2$$

Taylor's series is :

$$y = y_0 + h.y'_0 + \frac{h^2}{2!} y''_0 + \frac{h^3}{3!} y'''_0 + \dots$$

$$\therefore y(0.1) = 1 + 0.1(-1) + 0 + \frac{(0.1)^3}{3!} \quad (2)$$

$$\therefore y(0.1) = 0.9003$$

- (c) Find the value of the integral $\int_0^1 \frac{x^2}{1+x^3} dx$ using (i) Trapezoidal rule (ii) Simpson's $(1/3)^{rd}$ rule (iii) Simpson's $(3/8)^{th}$ rule. [8]

Ans: Let $I = \int_0^1 \frac{x^2}{1+x^3} dx$

$a=0, b=1$

Dividing limits into 4 parts i.e $n=4 \quad \therefore h = \frac{b-a}{n} = \frac{1}{4} = 0.25$

$x_0 = 0$	$x_1 = 0.25$	$x_2 = 0.50$	$x_3 = 0.75$	$x_4 = 1.0$
$y_0 = 0$	$y_1 = 0.06153$	$y_2 = 0.2222$	$y_3 = 0.39560$	$y_4 = 0.5$

(i) Trapezoidal rule: $I = \frac{h}{2} [X + 2R]$ -----(1)

$X = \text{sum of extreme ordinates} = y_0 + y_4 = 0 + 0.5 = 0.5$

$R = \text{sum of remaining ordinates} = y_1 + y_2 + y_3$
 $= 0.06153 + 0.2222 + 0.39560 = 0.67933$

$I = \frac{0.25}{2} (0.5 + 2(0.67933))$ (from 1)

$$\therefore I = 0.2323$$

(ii) Simpson's $(1/3)^{rd}$ rule:

$I = \frac{h}{3} [X + 2E + 4O]$ -----(2)

$X = \text{sum of extreme ordinates} = y_0 + y_4 = 0 + 0.5 = 0.5$

$E = \text{sum of even base ordinates} = y_2 = 0.2222$

$O = \text{sum of odd base ordinates} = y_1 + y_3 = 0.06153 + 0.39560 = 0.45713$

$I = \frac{0.25}{3} (0.5 + 2 \times 0.2222 + 4 \times 0.45713)$ (from 2)

$$\therefore I = 0.23108$$

(iii) Simpson's (3/8)th rule :

$$I = \frac{3h}{8} [X + 2T + 3R] \quad \text{-----(3)}$$

$$X = \text{sum of extreme ordinates} = y_0 + y_4 = 0 + 0.5 = 0.5$$

$$T = \text{sum of multiple of three base ordinates} = y_3 = 0.39560$$

$$R = \text{sum of remaining ordinates} = y_1 + y_2 = 0.06153 + 0.2222 = 0.28373$$

$$I = \frac{3 \times 0.25}{8} (0.5 + 2 \times 0.39560 + 3 \times 0.28373)$$

$$\therefore I = 0.2008$$

Q. 5 (a). Solve $(y - xy^2)dx - (x + x^2y)dy = 0$ [6]

Ans : $(y - xy^2)dx - (x + x^2y)dy = 0$ -----(1)

Comparing the given eqn with $M dx + N dy = 0$

$$\therefore M = (y - xy^2) \quad \therefore N = -(x + x^2y)$$

$$\frac{\partial M}{\partial y} = 1 - 2xy \quad \frac{\partial N}{\partial x} = -(1 + 2xy)$$

$$\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$$

The given differential eqn is not exact diff. eqn.

But the given diff. eqn is in the form of $y \cdot f(xy)dx + x f(xy)dy = 0$

$$\text{Integrating factor} = \text{I.F.} = \frac{1}{Mx - Ny} = \frac{1}{xy - x^2y^2 + xy + x^2y^2} = \frac{1}{2xy}$$

Multiply the I.F. to eqn (1)

$$\left(\frac{1}{2x} - \frac{y}{2}\right) dx - \left(\frac{1}{2y} + \frac{x}{2}\right) dy = 0$$

$$\therefore M_1 = \left(\frac{1}{2x} - \frac{y}{2}\right) \quad N_1 = -\left(\frac{1}{2y} + \frac{x}{2}\right)$$

$$\int M_1 dx = \int \left(\frac{1}{2x} - \frac{y}{2}\right) dx = \frac{1}{2} (\log x) - \frac{xy}{2}$$

$$\frac{\partial}{\partial y} \int M_1 dx = \frac{-x}{2}$$

$$\int [N_1 - \frac{\partial}{\partial y} \int M_1 dx] dy = \int \frac{-1}{2y} dy = \frac{-1}{2} (\log y)$$

The solution of given diff. eqn is given by,

$$\int M_1 dx + \int \left[N_1 - \frac{\partial}{\partial y} \int M_1 dx \right] dy = c$$

$$\therefore \frac{1}{2}(\log x) - \frac{xy}{2} - \frac{1}{2}(\log y) = c$$


$$\therefore \log \left(\frac{\sqrt{x}}{\sqrt{y}} \right) - \frac{xy}{2} = c$$

(b) Evaluate $\iiint \sqrt{1 - \frac{x^2}{a^2} - \frac{y^2}{b^2} - \frac{z^2}{c^2}} dx dy dz$ over the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$.

$$\frac{z^2}{c^2} = 1.$$

[8]

Ans : Ellipsoid : $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2}$



Cartesian coordinates \longrightarrow spherical coordinate system

$(x, y, z) \longrightarrow (r, \theta, \phi)$

Put $x = a r \sin \theta \cos \phi$, $y = b r \sin \theta \sin \phi$, $z = c r \cos \theta$

$$dx dy dz = abc r^2 \sin \theta dr d\theta d\phi$$

$$\therefore \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = r^2$$

$$f(x, y, z) = \sqrt{1 - \frac{x^2}{a^2} - \frac{y^2}{b^2} - \frac{z^2}{c^2}} = \sqrt{1 - r^2} = f(r, \theta, \phi)$$

Limits : $0 \leq r \leq 1$

$$0 \leq \theta \leq \frac{\pi}{2}$$

$$0 \leq \phi \leq \frac{\pi}{2}$$

$$\begin{aligned}
 \therefore I &= 8 \int \int \int \sqrt{1-r^2} abc r^2 \sin \theta dr d\theta d\phi \\
 &= 8 \int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \int_0^r \sqrt{1-r^2} abc r^2 \sin \theta dr d\theta d\phi \\
 &= 8 abc \int_0^{\frac{\pi}{2}} \sin \theta d\theta \int_0^{\frac{\pi}{2}} d\phi \int_0^r \sqrt{1-r^2} r^2 dr \\
 &= 8 abc [-\cos \theta]_0^{\frac{\pi}{2}} [\phi]_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \cos t \cdot \sin^2 t \cdot \cos t dt \quad \text{-----}\{ \text{put } r = \sin t \} \\
 &= 8 abc \left(\frac{\pi}{2}\right) \left(\frac{\pi}{8}\right) \quad \text{-----}\{ \text{use } \beta f^n \}
 \end{aligned}$$

$$\therefore I = \frac{\pi^2}{4} (abc)$$

(c) Evaluate $(2x + 1)^2 \frac{d^2y}{dx^2} - 2(2x + 1) \frac{dy}{dx} - 12y = 6x$ [8]

Ans : $(2x + 1)^2 \frac{d^2y}{dx^2} - 2(2x + 1) \frac{dy}{dx} - 12y = 6x$ (1)

Put $(2x + 1) = e^z \quad \Rightarrow \quad x = \frac{e^z - 1}{2}$

$\frac{dz}{dx} = \frac{2}{(2x+1)}$ but $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx} = 2 \frac{dy}{dz} = \frac{2}{(2x+1)} Dy$ where $D = \frac{d}{dz}$

$\therefore (2x+1) \frac{dy}{dx} = 2 Dy$

$\therefore (2x + 1)^2 \frac{d^2y}{dx^2} = 2^2 D(D - 1)y$

From (1),

$4D(D - 1)y - 4 Dy - 12y = 6\left(\frac{e^z - 1}{2}\right)$

$(4D^2 - 8D - 12)y = 3(e^z - 1)$

For complementary solution ,

$(4D^2 - 8D - 12) = 0$

$\therefore D = -1, 3$

$$\therefore y_c = c_1 e^{-z} + c_2 e^{3z}$$

For particular integral ,

$y_p = \frac{1}{f(D)} X$

$$y_p = \frac{1}{4D^2 - 8D - 12} (3(e^z - 1))$$

$$\therefore y_p = \frac{3}{4} \frac{1}{D^2 - 2D - 3} (e^z - 1) \quad \text{put } D = a = 1 \text{ and } D = a = 0$$

$$\therefore y_p = \frac{3}{4} \left(\frac{1}{3} - \frac{e^z}{4} \right)$$

The general solution of given differential eqn is ,

$$\therefore y_g = y_c + y_p = c_1 e^{-z} + c_2 e^{3z} + \frac{3}{4} \left(\frac{1}{3} - \frac{e^z}{4} \right)$$

Resubstituting z ,

$$\therefore y_g = c_1 (2x + 1)^{-1} + c_2 (2x + 1)^3 + \frac{3}{4} \left(\frac{1}{3} - \frac{(2x+1)}{4} \right)$$

Q.6.(a) A resistance of 100 ohms and inductance of 0.5 henries are connected in series With a battery of 20 volts. Find the current at any instant if

the relation between L, R, E is $L \frac{di}{dt} + Ri = E$. [6]

Ans : $L \frac{di}{dt} + Ri = E$

$$\therefore \frac{di}{dt} + \frac{Ri}{L} = \frac{E}{L}$$

Solution is given by ,

$$i \cdot e^{\int \left(\frac{R}{L}\right) dt} = \int e^{\int \left(\frac{R}{L}\right) dt} \cdot \frac{E}{L} dt + c$$

$$\therefore i \cdot e^{(Rt/L)} = \frac{E e^{(Rt/L)}}{R} + c$$

At $t=0, i=0 \quad \therefore c = -\frac{E}{R}$

$$\therefore i \cdot e^{(Rt/L)} = \frac{E e^{(Rt/L)}}{R} - \frac{E}{R}$$

$$\therefore i = \frac{E}{R} (1 - e^{-(Rt/L)})$$

For given condition $R = 100, L = 0.5, E = 20$

$$\therefore i = 0.2(1 - e^{-200t})$$

(b) Solve by variation of parameter method $\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + 2y = e^{ex}$.

[6]

Ans : $\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + 2y = e^{ex}$

Put $D = \frac{d}{dx}$ $\therefore D^2y + 3Dy + 2y = 0$

For complementary solution,

$$f(D)=0$$

$$\therefore D^2 + 3D + 2 = 0$$

$$D = -1, -2$$

$$\therefore y_c = c_1e^{-x} + c_2e^{-2x}$$

Particular integral is given by ,

$$y_p = y_1p_1 + y_2p_2$$

where $p_1 = \int \frac{-y_2X}{w} dx$

$$p_2 = \int \frac{y_1X}{w} dx$$

$$w = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$$

$$\therefore w = \begin{vmatrix} e^{-x} & e^{-2x} \\ -e^{-x} & -2e^{-2x} \end{vmatrix} = -e^{-3x}$$

$$p_1 = \int \frac{e^{-2x} \cdot e^{ex}}{e^{-3x}} dx = \int e^{ex} \cdot e^x dx = \int e^t dt = e^t \dots \dots \{ \text{put } e^x = t \Rightarrow e^x dx = dt \}$$

$$p_2 = \int \frac{e^{-x}}{-e^{-3x}} \cdot e^{ex} dx = \int e^{ex} \cdot e^{2x} dx = \int t \cdot e^t dt = e^t x - e^t$$

$$\therefore y_p = e^x e^{ex} - (e^x e^{ex} - e^{ex}) \cdot e^{-2x} = e^{-2x} \cdot e^{ex}$$

The general solution of given differential eqn is given by ,

$$y_g = y_c + y_p = c_1e^{-x} + c_2e^{-2x} + e^{-2x} \cdot e^{ex}$$


(c) Evaluate $\int \int xy(x - 1) dx dy$ over the region bounded by $xy = 4, y = 0, x = 1$ and $x = 4$ [8]

Ans : Let $I = \int \int xy(x - 1) dx dy$

Rectangular hyperbola : $xy = 4$ Lines : $x = 1, x = 4, y = 0$

Intersection of line $x = 1$ and $xy = 4$ is $(1,4)$.

Intersection of line $x = 4$ and $xy = 4$ is $(4,1)$



$$\therefore \quad 0 \leq y \leq \frac{x}{4}$$
$$1 \leq x \leq 4$$

$$\begin{aligned} \therefore I &= \int_1^4 \int_0^{\frac{x}{4}} (x^2 y - xy) dy dx \\ &= \int_1^4 \left[\frac{y^2}{2} x^2 - \frac{y^2 x}{2} \right]_0^{\frac{x}{4}} dx \\ &= \int_1^4 \left(8 - \frac{8}{x} \right) dx \\ &= [8x - 8 \log x]_1^4 \end{aligned}$$

$$\therefore I = 8(3 - 2 \log 2)$$